



# IMMUNOSUPPRESSION INDUCED BY CANCER CELLS: PD-1 AND CTLA-4 PATHWAYS

## INTRODUCTION


- o Cancer cells induce **immunologic tolerance** → absence of an immune response against certain antigens, causing the escape of these cells from the immune system.
- o 2 groups of tumor **antigens**: **true tumor-specific antigens** (encoded by mutant cellular genes), and tumor-associated antigens (encoded by normal cellular genes).



**Figure 1. Immunosuppressive strategies.** (1) impairment of the Ag presentation/processing machinery, (2) defects in TCR proximal signals, (3) secretion of immunoregulatory cytokines, (4) negative costimulatory signals, (5) tryptophan depletion, (6) proapoptotic pathways, and (7) regulatory T-cells. Adapted from reference [2].

## PD-1 PATHWAY

- ❖ Programmed cell death-1 is a 288 aa protein that belongs to Ig superfamily.
- Consists in an extracellular IgV-like domain, and a cytoplasmic region (with an ITIM and an ITSM). — regulator of T-cell function (figure 2).
- ❖ Ligands: PD-L1 (B7-H1) and PD-L2 (B7-DC), expressed in lymphoid, in non-lymphoid cells and in tumor cells.




**Figure 3. PD-1 signaling.** (1) T-cell activation, (2) PD-1 expression, (3) PD-1/PD-L interaction, (4) ITSM phosphorylation, (5) SHP-2 recruitment and phosphorylation, (6+7) dephosphorylation of proximal and downstream effector molecules. Adapted from reference [10].

\*It has been observed in PD-1 KO mice the loss of peripheral tolerance and the development of autoimmunity.

Apoptosis, energy, exhaustion, molecular shield, IL-10 production → **T-cell tolerance**

## CTLA-4 PATHWAY

- ❖ CTLA-4 (Cytotoxic T-lymphocyte associated antigen protein 4), is a 223 aa glycoprotein that belongs to Ig superfamily and has an extracellular IgV-like domain. — regulator of T-cell activation (figure 4).
- ❖ Ligands: B7-1 (CD80) and B7-2 (CD86), expressed in APCs and in tumor cells.




**Figure 4. Solution structure of human CTLA-4 with a tetrasaccharide core attached.** Reference [26].

\*CTLA-4 KO mice developed autoimmune diseases and died at 3-4 weeks old → significant role in the development of peripheral tolerance to self-proteins.

## DIFFERENCES BETWEEN PD-1 AND CTLA-4

|                                    | CTLA-4     | PD-1                               | Implications   |
|------------------------------------|------------|------------------------------------|--|
| Expression                         | T-cells    | T-cells, B-cells and myeloid cells | PD-1: more broadly role in the regulation of immune responses  |
| Interaction with AP2 (endocytosis) | Yes        | No                                 | CTLA-4 undetectable on the cell surface; PD-1 greater expression   |
| SHP-2 association                  | Indirectly | Directly                           | CTLA-4 preserves a little PI3K activity, whereas PD-1 affects a more global inhibition of T-lymphocytes (figure 6) |



**Figure 6. T-lymphocyte inhibition by CTLA-4 and PD-1.** CTLA-4 preserves some PI3K activity but inhibits Akt directly. PD-1 inhibits PI3K directly. Adapted from reference [35].

## IMPLICATIONS FOR CANCER IMMUNOTHERAPY

A variety of checkpoint blocking agents have been developed to block PD-1 and CTLA-4 signaling (including monoclonal Ab). Examples: Nivolumab and Ipilimumab (anti-PD-1), Ipilimumab (anti-CTLA-4).

### PROS

- ✓ More effective than classical therapies in metastatic cancers.
- ✓ Synergistic activity → combinatorial therapies.

### CONS

- ✗ Side effects.
- ✗ Very new, we don't know the long-term side effects.
- ✗ Currently very expensive therapies → not accessible to everyone.

## CONCLUDING REMARKS

**1** PD-1 and CTLA-4: powerful negative regulators of the immune response (KO experiments, development of autoimmunity) → very important role in the regulation of the IS, immunosuppression.

**2** Cancer uses this mechanism to evade and escape from IS cells → tolerance toward cancer cells, leading to the development of malignant tumors.

**3** Therapy: it's difficult to find a monotherapy that works at 100% because there are many different molecules and mechanisms involved, → combinatorial therapies should be tested.

## REFERENCES

- [1] Magera, M.Y. and Sykes, M., *Journal of Clinical Oncology*, 2004, 22: 1136-1151. [2] Rabinovich, G.A., Gabrilovich, D., and Sotomayor E.M., *Annual Review of Immunology*, 2007, 25: 267-296. [3] Zou, W., and Chen, L., *Nature Reviews Immunology*, 2008, 8: 467-477. [4] Ishida, Y., *The EMBO Journal*, 1992, 11: 3887-3895. [5] Okazaki, T., and Honjo, T., *Trends in Immunology*, 2006, 27: 195-201. [6] NCBI (2014) PDCD1 gene. URL: <http://www.ncbi.nlm.nih.gov/ncbi/podcasts/2014> [consulted: 26-03-2014]. [7] Uniprot (2014) Programmed cell death protein 1. URL: <http://www.uniprot.org/uniprot/P13113> [consulted: 26-03-2014]. [8] Freeman, G.J. et al., *The Journal of Experimental Medicine*, 2000, 192: 1027-1034. [9] Neri, L. et al., *J. Immunol.*, 2006, 177: 5028-5035. [10] Latchman, Y. et al., *Proc. Natl. Acad. Sci. USA*, 2002, 99: 12293-12297. [11] Blank, C. et al., *Int. J. Cancer*, 2006, 119: 317-321. [12] Nishimura, H. et al., *Science*, 2001, 291: 319-322. [13] eBioScience (2012) CTLA4 signaling. URL: [http://www.ebioscience.com/resources/pathways/ctla4\\_signaling\\_pathway.htm](http://www.ebioscience.com/resources/pathways/ctla4_signaling_pathway.htm) [consulted: 26-03-2014]. [14] NCBI (2014) CTLA4 gene. URL: <http://www.ncbi.nlm.nih.gov/ncbi/podcasts/2014> [consulted: 26-03-2014]. [15] Uniprot (2014) Cytotoxic T-lymphocyte protein 4. URL: <http://www.uniprot.org/uniprot/P13113> [consulted: 26-03-2014]. [16] Protein Data Bank (2014) CTLA-4. URL: <http://www.rcsb.org/pdb/explore/explorer/structure/doi/10.1093/emboj/2014/2501311> [consulted: 26-03-2014]. [17] Jago, C. B. et al., *Clin. Exp. Immunol.*, 2004, 136: 463-471. [18] Nirschl, C.J., and Drake C.G., *Clinical Cancer Research*, 2013, 19: 4917-4924. [19] Bhatia, S. et al., *Immunology Letters*, 2006, 104: 70-75. [20] Chuang, E. et al., *Immunology*, 2000, 13: 313-322. [21] Lee, K. M. et al., *Science*, 1998, 282: 2263-2266. [22] Wilunas, T. L., Bakker, C. Y., and Bluestone, J. A., *J. Exp. Med.*, 1996, 183: 2541-2550. [23] Salomon, B., and Bluestone, J.A., *Annual Review of Immunology*, 2001, 19: 225-252. [24] Tholi, E. A. et al., *Immunology*, 1995, 3: 541-547. [25] Parry, R. V. et al., *Molecular and Cellular Biology*, 2005, 25: 9543-9553. [26] Wolchok, I. D. et al., *The New England Journal of Medicine*, 2013, 369: 122-133. [27] Ornd Hamid, M. D. et al., *The New England Journal of Medicine*, 2013, 369: 134-144. [28] Curran, M. A. et al., *PNAS*, 2010, 107: 4275-4280. [29] Alegre, M.L., Frauwirth, K.A., and Thompson, C.B., *Nature Reviews Immunology*, 2001, 1: 220-228.